

2. Burgess LJ, Reuter H, Carstens ME, et al. The use of adenosine deaminase and interferon-gamma as diagnostic tools for tuberculous pericarditis. *Chest* 2002;122:900-5.
3. Mayosi BM, Burgess LJ, Doubell AF. Tuberculous pericarditis. *Circulation* 2005;112:3608-16.
4. Fowler NO. Tuberculous pericarditis. *JAMA* 1991;266:99-103.
5. Lee JH, Lee CW, Lee SG, et al. Comparison of polymerase chain reaction with adenosine deaminase activity in pericardial fluid for the diagnosis of tuberculous pericarditis. *Am J Med* 2002;113:519-21.

Circadian Variations of Ventricular Arrhythmias and Sleep-Disordered Breathing in HF Patients



We read with interest the study by Patton et al. (1) on the unexpected absence of typical circadian variation of ventricular arrhythmias observed in the SCD-HeFT (Sudden Cardiac Death in Heart Failure Trial). Sleep-disordered breathing (SDB), broadly categorized into obstructive and central sleep apnea, has been associated with increased cardiovascular morbidity and mortality. Cardiac arrhythmias are responsible for some of the higher cardiovascular morbidity and mortality rates observed in patients with SDB. The association between atrial fibrillation and SDB is well established, although the association between SDB and life-threatening ventricular arrhythmias also seems plausible. Obstructive sleep apnea predicts sudden cardiac death independently of other well-established risk factors (2), and, unlike the general population, patients with SDB have a higher incidence of sudden cardiac death during sleep (3). Studies consistently report an SDB prevalence of $\geq 50\%$ in the chronic heart failure population. The prevalence of SDB in patients with an implantable cardioverter-defibrillator (ICD) ranges between 57.8% and 66.3% (4,5). In a cohort of 472 ICD patients with heart failure receiving cardiac resynchronization therapy, a significant risk enhancement of ventricular arrhythmias and appropriate ICD therapies owing to both central and obstructive sleep apnea was found (5). Importantly, for heart failure patients with a primary inappropriate ICD therapies (4). Patton et al. (1) observed an increase in the onset of ventricular arrhythmias during sleep in patients with an ICD and SDB. Data on SDB for patients enrolled in the SCD-HeFT were not reported. Thus, it is our opinion that the observed deviation in circadian variation of ventricular arrhythmias reported by Patton et al. (1) may be influenced, at

least in part by the presence of SDB, a very prevalent condition among heart failure patients with an ICD.

*Miguel A. Arias, MD, PhD

Marta Pachón, MD

Finn Akerström, MBChB

Alberto Puchol, MD

Luis Rodríguez-Padial, MD, PhD

*Unidad de Arritmias y Electrofisiología Cardíaca

Hospital Virgen de la Salud

Avda. Barber 30

45004 Toledo

Spain

E-mail: maapalomares@secardiologia.es

<http://dx.doi.org/10.1016/j.jacc.2014.07.984>

REFERENCES

1. Patton KK, Hellkamp AS, Lee KL, et al. Unexpected deviation in circadian variation of ventricular arrhythmias: the SCD-HeFT (Sudden Cardiac Death in Heart Failure Trial). *J Am Coll Cardiol* 2014;63:2702-8.
2. Gami AS, Olson EJ, Shen WK, et al. Obstructive sleep apnea and the risk of sudden cardiac death: a longitudinal study of 10,701 adults. *J Am Coll Cardiol* 2013;62:610-6.
3. Gami AS, Howard DE, Olson EJ, Somers VK. Day-night pattern of sudden death in obstructive sleep apnea. *N Engl J Med* 2005;352:1206-14.
4. Kreuz J, Skowasch D, Horlbeck F, et al. Usefulness of sleep-disordered breathing to predict occurrence of appropriate and inappropriate implantable-cardioverter defibrillator therapy in patients with implantable cardioverter-defibrillator for primary prevention of sudden cardiac death. *Am J Cardiol* 2013;111:1319-23.
5. Bitter T, Westerheide N, Prinz C, et al. Cheyne-Stokes respiration and obstructive sleep apnoea are independent risk factors for malignant ventricular arrhythmias requiring appropriate cardioverter-defibrillator therapies in patients with congestive heart failure. *Eur Heart J* 2011;32:61-74.

REPLY: Circadian Variations of Ventricular Arrhythmias and Sleep-Disordered Breathing in HF Patients



We appreciate the interest of Dr. Arias and colleagues in our study of circadian and septadian patterns of implantable cardioverter-defibrillator therapy in the SCD-HeFT (Sudden Cardiac Death in Heart Failure Trial) population (1). In their Letter, they relevantly highlight the importance of sleep-disordered breathing as a trigger of ventricular arrhythmias and implantable cardioverter-defibrillator therapies. Both central and obstructive sleep apnea exert strong effects on the autonomic nervous system and are known to be proarrhythmic (2).

We agree with Dr. Arias and colleagues that sleep-disordered breathing is an important and increasingly recognized trigger of arrhythmias (3). Unfortunately, we do not have information on the presence